
Usmle Step 2 Ck Sample Questions

*Usmle Step 2 Ck
Sample Questions*

*Downloaded from
old.oplm.com by
Chambers & Willis*

MASTERING THE USMLE STEP 2 CK: THE POWER OF PRACTICE QUESTIONS

Every medical student knows the feeling. You have spent years absorbing textbooks, lecture notes, and clinical experiences. Now, one exam stands between you and your residency application: the USMLE Step 2 CK. This test does not just measure your knowledge; it measures your ability to apply that knowledge under pressure. The single most effective way to prepare is to work through realistic **Usmle Step 2 Ck Sample Questions** that mirror the style, difficulty, and clinical reasoning demands of the actual exam. This article will explore why sample questions are indispensable, walk you through several detailed examples, and provide strategies to maximize your performance on test day.

WHY USMLE STEP 2 CK SAMPLE QUESTIONS ARE ESSENTIAL

Reading textbooks and watching lecture videos builds a foundation, but the Step 2 CK is fundamentally a test of clinical decision-making. You are not simply asked to recall a fact. Instead, you are presented with a patient vignette and must synthesize information, prioritize diagnoses, select the next best step, or choose the most appropriate management. Practicing with high-quality sample questions trains your

brain to think in this specific way. It forces you to integrate internal medicine, surgery, pediatrics, obstetrics and gynecology, psychiatry, and preventive medicine into a single, coherent thought process.

Furthermore, exposure to sample questions helps you identify patterns. Certain presentations recur frequently, and recognizing these patterns under timed conditions is a skill that must be cultivated. Many students find that their score improves dramatically simply by becoming familiar with the question format and the nuances of how answer choices are constructed. This is why dedicated practice with **Usmle Step 2 Ck Sample Questions** is not merely helpful; it is a cornerstone of any serious study plan.

UNDERSTANDING THE STRUCTURE OF STEP 2 CK QUESTIONS

Before diving into examples, it helps to understand the anatomy of a typical Step 2 CK question. Each question begins with a clinical vignette. This vignette provides a patient scenario including age, gender, presenting complaint, history of present illness, past medical history, medications, physical examination findings, and often laboratory or imaging results. The vignette is followed by a single best answer question, with five or more answer choices. There are no multiple correct answers, no "select all that apply" formats. Your task is to choose the single best option.

The questions are designed to test your ability to:

- Interpret clinical data accurately
- Generate a differential diagnosis
- Select the most appropriate diagnostic test
- Choose the correct initial management or treatment
- Recognize when to refer or admit a patient
- Apply preventive medicine and screening guidelines

Working through a large volume of **Usmle Step 2 Ck Sample Questions** is the best way to sharpen each of these skills.

SAMPLE QUESTION 1: **CARDIOLOGY**

Let us begin with a classic cardiology scenario. This type of question tests your ability to recognize an acute coronary syndrome and select appropriate management.

Vignette

A 62-year-old man with a history of hypertension and hyperlipidemia presents to the emergency department with substernal chest pressure that began two hours ago. The pain radiates to his left arm and is associated with diaphoresis and nausea. His blood pressure is 145/90 mm Hg, heart rate is 102 bpm, and respiratory rate is 18 breaths per minute. An electrocardiogram reveals ST-segment elevation in leads V2, V3, and V4. Cardiac troponin is elevated. The patient has no contraindications to anticoagulation.

Question

Which of the following is the most appropriate next step in management?

1. Administer tissue plasminogen activator (tPA)
2. Start heparin and schedule stress test in 48 hours
3. Transfer for emergent percutaneous coronary intervention (PCI)
4. Start medical management with beta-blockers and nitrates alone
5. Give aspirin and discharge with follow-up

Answer and Explanation

The correct answer is option 3: Transfer for emergent percutaneous coronary intervention (PCI). This patient presents with ST-segment elevation myocardial infarction (STEMI). The standard of care for STEMI within 12 hours of symptom onset is emergent reperfusion, preferably via PCI within 90 minutes of arrival. While tPA is an option if PCI is not available, guidelines favor PCI when it can be performed in a timely manner. Simply starting heparin or managing medically without reperfusion is insufficient. This question underscores a key theme in **Usmle Step 2 Ck Sample Questions**: you must know the most current, evidence-based guidelines for acute management.

SAMPLE QUESTION 2: **PEDIATRICS**

Pediatric questions often revolve around developmental milestones, common

childhood illnesses, and growth parameters. This example focuses on failure to thrive.

Vignette

A 14-month-old girl is brought to the clinic by her mother for a routine well-child visit. The mother reports that the child eats poorly and is difficult to feed. The child was born at full term with a normal birth weight. On examination, the child is irritable and appears thin. Weight is at the 3rd percentile, height is at the 10th percentile, and head circumference is at the 25th percentile. There is noticeable loss of subcutaneous fat. The remainder of the physical examination is unremarkable.

Question

Which of the following is the most likely cause of this child's condition?

1. Gastroesophageal reflux disease
2. Inadequate caloric intake
3. Cystic fibrosis
4. Celiac disease
5. Growth hormone deficiency

Answer and Explanation

The correct answer is option 2: Inadequate caloric intake. This child presents with failure to thrive, characterized by weight falling below the 3rd percentile with relative preservation of height and head circumference. This pattern suggests a caloric deficit rather than an underlying endocrine or organic disorder. Inadequate intake is the most common cause of failure to thrive in this

age group, often due to behavioral feeding difficulties or improper feeding practices. While cystic fibrosis and celiac disease can cause malabsorption and poor growth, they typically present with additional signs such as steatorrhea, respiratory symptoms, or abdominal distension. Growth hormone deficiency would more significantly affect height. Reflux is common but rarely causes such profound growth failure in isolation. This question highlights the importance of distinguishing between organic and non-organic causes of growth failure, a recurring theme in **Usmle Step 2 Ck Sample Questions**.

SAMPLE QUESTION 3: OBSTETRICS AND GYNECOLOGY

OB-GYN questions often involve pregnancy complications, abnormal uterine bleeding, or gynecologic infections. This example deals with postpartum complications.

Vignette

A 30-year-old woman, gravida 2 para 2, presents to the emergency department five days after a spontaneous vaginal delivery. She complains of heavy vaginal bleeding, fever, and lower abdominal pain. Her temperature is 38.9°C (102°F), heart rate is 110 bpm, and blood pressure is 100/60 mm Hg. On bimanual examination, the uterus is tender, enlarged, and boggy. There is foul-smelling lochia. Laboratory studies show leukocytosis with a left shift.

Question

Which of the following is the most appropriate next step in management?

1. Perform dilation and curettage

- (D&C) for retained products
- 2. Administer broad-spectrum intravenous antibiotics
- 3. Start oxytocin infusion
- 4. Obtain a pelvic ultrasound
- 5. Administer methylergonovine

Answer and Explanation

The correct answer is option 2: Administer broad-spectrum intravenous antibiotics. This patient presents with signs and symptoms consistent with postpartum endometritis. Endometritis is an infection of the endometrial lining that typically occurs after delivery, especially in the setting of prolonged labor, chorioamnionitis, or cesarean section. The combination of fever, uterine tenderness, and foul-smelling lochia is highly suggestive. The first step in management is to initiate broad-spectrum antibiotics covering gram-positive, gram-negative, and anaerobic organisms. While a pelvic ultrasound can help evaluate for retained products of conception, it should not delay antibiotics in an unstable patient. D&C may be indicated if retained products are found, but it is not the first step. Oxytocin and methylergonovine are used for postpartum hemorrhage secondary to uterine atony, not for infection. This question teaches an important principle: when infection is suspected, antibiotics come first. Working through **Usmle Step 2 Ck Sample Questions** like this one helps solidify your approach to postpartum emergencies.

SAMPLE QUESTION 4: **PSYCHIATRY**

Psychiatry questions often test your

ability to diagnose based on DSM-5 criteria and select appropriate pharmacotherapy or psychotherapy.

Vignette

A 28-year-old woman presents to the outpatient clinic with complaints of persistent sadness, low energy, and difficulty sleeping for the past three weeks. She reports that her symptoms began shortly after she lost her job. She has lost her appetite and feels worthless. She denies any suicidal ideation or psychotic symptoms. She has no prior history of mood disorders. Her medical history is unremarkable, and she takes no medications.

Question

Which of the following is the most appropriate initial treatment for this patient?

- 1. Start a selective serotonin reuptake inhibitor (SSRI)
- 2. Refer for cognitive behavioral therapy (CBT)
- 3. Hospitalize for safety monitoring
- 4. Start a benzodiazepine for sleep
- 5. Recommend exercise and watchful waiting for two more weeks

Answer and Explanation

The correct answer is option 2: Refer for cognitive behavioral therapy (CBT). This patient meets criteria for a major depressive episode, but the duration is only three weeks. According to DSM-5, the symptoms must be present for at least two weeks, so she does meet

duration criteria. However, for mild to moderate depression, especially with a clear precipitant like job loss, psychotherapy is recommended as first-line treatment before initiating medication. She does not endorse suicidal ideation, so hospitalization is not required. While SSRIs are effective, guidelines suggest trying CBT first for mild cases. Benzodiazepines are not recommended for the treatment of depression. Watchful waiting may be appropriate for very mild symptoms, but this patient's symptoms are already impacting her function significantly. This question emphasizes the stepped-care approach to mental health treatment, a concept you will see repeated in **Usmle Step 2 Ck Sample Questions**.

STRATEGIES FOR ANSWERING STEP 2 CK QUESTIONS

Now that you have seen several example questions, let us discuss strategies for tackling them effectively. These techniques will serve you well across all subject areas.

Read the Question First

Many test-takers recommend reading the question stem before reading the vignette. This allows you to know what you are looking for as you read the clinical data. For example, if the question asks for the "most likely diagnosis," you will focus on fitting the findings into a diagnostic framework. If it asks for the "next best step," you will pay closer

attention to what has already been done. This small adjustment can save time and reduce cognitive overload.

Identify Key Details

Not every detail in a vignette is equally important. Train yourself to highlight the critical pieces: age, gender, presenting complaint, duration of symptoms, vital signs, and key physical findings. Laboratory and imaging results often confirm or rule out possibilities in your differential. When practicing with **Usmle Step 2 Ck Sample Questions**, get into the habit of mentally summarizing the case in one or two sentences before looking at the answer choices.

Generate a Differential Before Reading Answers

Cover the answer choices with your hand or mentally block them out. Based on the vignette alone, generate your own list of possible diagnoses. Then, uncover the answers and see which one matches your thinking. This prevents you from being swayed by plausible-sounding wrong answers. The USMLE is notorious for including answer choices that are partially correct but not the best option.

A