
How To Get Your Period Back

How To Get Your Period Back Downloaded from
old.oplm.com by Guerra
& Isaiah

UNDERSTANDING MENSTRUAL HEALTH: A NATURAL GUIDE

Losing your period, a condition medically known as amenorrhea, can be concerning whether you are trying to conceive, managing a health condition, or simply want your body to return to its natural rhythm. Your menstrual cycle is a vital sign of overall health, reflecting hormonal balance, nutritional status, and stress levels. If you are wondering **how**

to get your period back, it is essential to approach the process holistically, addressing underlying causes rather than forcing a cycle. This guide provides evidence-based strategies to help restore regular ovulation and menstruation naturally.

WHY DID YOU LOSE YOUR PERIOD? COMMON UNDERLYING CAUSES

Before exploring solutions, it is critical to identify why your period stopped. The most common reasons include:

- **Functional Hypothalamic Amenorrhea (FHA):** Often caused by chronic stress, excessive exercise, or undereating (low energy availability). The hypothalamus stops releasing gonadotropin-releasing hormone (GnRH), halting ovulation.
- **Polycystic Ovary Syndrome (PCOS):** A hormonal imbalance leading to irregular ovulation or anovulation due to elevated androgens or insulin resistance.
- **Low Body Weight or Eating Disorders:** Body fat below a critical threshold disrupts hormone production (e.g., leptin).
- **Hormonal Contraceptives:** Post-pill amenorrhea (*post-birth control syndrome*) can occur as the body

readjusts to its natural cycle.

- **Thyroid or Prolactin Disorders:** Thyroid imbalances or elevated prolactin can interfere with menstrual regulation.
- **Premature Ovarian Insufficiency (POI):** Ovaries stop functioning normally before age 40.

If you have not had a period for three months or more (or six months prior if cycles were irregular), consult a healthcare provider for blood work, imaging, and a thorough evaluation.

HOW TO GET YOUR PERIOD BACK: HOLISTIC STRATEGIES

1. Prioritize Calorie and Nutrient Sufficiency

Undereating is one of the most common drivers of amenorrhea. When your body perceives a calorie deficit, it conserves energy by shutting down non-essential functions like reproduction. **How to get your period back** often begins with eating more. Focus on:

- **Energy Availability:** Aim for at least 2,500–3,000 calories per day if you are active, adjusted for your

individual needs. Do not skip meals, and include carbohydrates, healthy fats, and protein at every meal.

- **Healthy Fats:** Hormones are built from cholesterol. Include full-fat dairy, avocados, nuts, seeds, olive oil, and fatty fish.
- **Complex Carbohydrates:** Whole grains, legumes, and root vegetables support stable blood sugar and leptin production.
- **Micronutrients:** Iron (red meat, spinach), zinc (pumpkin seeds, shellfish), magnesium (dark chocolate, leafy greens), and B vitamins (eggs, liver) are essential for ovulation.

If you have a history of restrictive eating

or an eating disorder, work with a registered dietitian specializing in menstrual health.

2. Reduce Physical Activity (Temporarily)

High-volume or high-intensity exercise combined with low calorie intake is a classic trigger for FHA. Even if you are eating normally, excessive training can raise cortisol and suppress GnRH. To restore your cycle:

- **Take a break:** Reduce exercise frequency and intensity by 30-50% for at least 2-4 weeks. Replace

heavy cardio with walking, gentle yoga, or restorative pilates.

- **Add rest days:** Schedule at least two full rest days per week and prioritize sleep (7-9 hours).
- **Listen to your body:** If you feel exhausted or notice poor recovery, your training load is too high.

For athletes under a coach, communicate your goal of restoring menstruation; many can adjust programming to support reproductive health.

3. Manage

Chronic Stress

Cortisol, the primary stress hormone, acts as a direct antagonist to reproductive hormones. When cortisol is chronically elevated, your body prioritizes survival over fertility. Incorporate these evidence-based stress-reduction techniques:

- **Mindful breathing:** Practice 4-7-8 breathing (inhale for 4 counts, hold for 7, exhale for 8) several times daily.
- **Limit caffeine and stimulants:** These can raise cortisol. Switch to herbal teas (chamomile, spearmint, ashwagandha).
- **Establish a wind-down routine:** No screens one hour before bed,

low lighting, and gentle stretching.

- **Consider adaptogens:** Herbs like ashwagandha, rhodiola, and maca root can help normalize cortisol levels, though consult a healthcare professional before supplementing.

If your stress is due to life circumstances, therapy (Cognitive Behavioral Therapy or counseling) can be invaluable in restoring hormonal balance.

4. Restore Healthy Body

Weight and Composition

In cases of low body weight (BMI under 18.5) or rapid weight loss, regaining weight is often the single most effective strategy. Studies show that many women with FHA recover their periods after gaining 2-5% of body weight, particularly in fat mass. This does not require “reckless eating” but rather a consistent increase in caloric density. For some, reaching a body fat percentage around 22-25% is necessary for ovulation to resume.

For those with PCOS and overweight, losing 5-10% of body weight through a balanced diet (low-glycemic, anti-

inflammatory) can restore ovulation. The approach is opposite: reduce insulin resistance through moderate caloric restriction and increased fiber intake. However, always avoid crash diets, as rapid weight loss can worsen amenorrhea.

5. Consider Supplements and Herbal Support

While diet is foundational, certain supplements may support the return of your period:

- **Inositol (myo-inositol & D-chiro-inositol):** Particularly

beneficial for PCOS-related amenorrhea, as it improves insulin sensitivity and promotes ovulation.

- **Zinc:** Low levels are linked to menstrual irregularities. Dosages of 30-40 mg per day (short-term) can be helpful.
- **Magnesium:** Helps reduce stress, improve sleep, and support hormonal detoxification.
- **Vitamin D:** Deficiency is common in women with amenorrhea. Adequate levels (checked via blood) are essential for ovulation.
- **Chasteberry (Vitex):** May help regulate prolactin and progesterone, often used in post-pill amenorrhea or elevated prolactin.

Always discuss supplements with a doctor, as some can interfere with medications or thyroid function.

MEDICAL INTERVENTIONS WHEN NATURAL APPROACHES ARE NOT ENOUGH

If lifestyle changes do not restore your period within 3-6 months, or if you have an underlying medical condition, medical help may be necessary.

Hormonal Testing and

Diagnosis

A healthcare provider can order:

- Blood tests: LH, FSH, estradiol, progesterone, prolactin, TSH, free T3, free T4, and androgens (testosterone, DHEA-S).
- Pelvic ultrasound to check for ovarian cysts (PCOS) or uterine issues.

used when trying to conceive.

- **Progesterone or Combination Birth Control:** Progesterone therapy can trigger withdrawal bleeding but does not address the root cause. It is sometimes used for a few months to “reset” cycles, but natural approaches are preferred.
- **Dopamine Agonists:** For elevated prolactin (e.g., cabergoline).

Medications

- **Metformin:** Often prescribed for PCOS to improve insulin sensitivity and induce ovulation.
- **Clomiphene or Letrozole:** Ovulation induction medications

When to See a Specialist

If you have not had a period for over a year, are under 40 and have premature

ovarian insufficiency, or if you experience symptoms like hot flashes or severe headaches, see a reproductive endocrinologist or gynecologist specialising in amenorrhea.

PATIENCE AND CONSISTENCY: THE TIMELINE FOR RECOVERY

Restoring your period is not an overnight process. The hypothalamus can take weeks to months to resume normal pulsatile GnRH secretion once the underlying stressor is removed. Some women see a period within 4-6 weeks of increasing energy intake and reducing exercise, while others may need 6-12 months of consistent healthy habits. Track your basal body temperature (BBT) or use ovulation predictor kits to detect a return of ovulation before actual

bleeding. If you notice fertile cervical mucus or a BBT shift, your body is on the right track.

It is common to experience missed periods intermittently during recovery. This does not mean you have failed – be patient with your body.

THE ROLE OF SLEEP AND CIRCADIAN RHYTHMS

Poor sleep disrupts the hypothalamic-pituitary-ovarian axis. Aim for 8 hours of quality sleep per night, ideally before midnight. Keep your bedroom cool and dark. If you have trouble sleeping, try magnesium glycinate or a warm Epsom salt bath. Circadian disruption from late-night screen time can raise evening cortisol, further suppressing LH.

CONCLUSION: HONOR YOUR BODY, FIND YOUR CYCLE

Learning **how to get your period back** is ultimately about listening to your body and addressing the root cause of the disruption. Whether it requires slowing down physically, eating more nutrient-dense food, managing stress, or seeking medical support for an underlying condition like PCOS or thyroid dysfunction, every step you take toward balance brings your cycle closer to

natural rhythm. Your period is not something to fear or suppress – it is a monthly sign of vitality. With patience, self-compassion, and evidence-based adjustments, many women are able to resume regular ovulation and menstruation. If progress stalls, professional guidance can make all the difference. Remember, true health is not measured by a single number on a scale, but by the harmony of your entire hormonal symphony.