
List Of Algebra 2 Formulas

*List Of
Algebra 2
Formulas*

*Downloaded
from
old.oplm.com
by Daniela &
Townsend*

INTRODUCTION: WHY A COMPREHENSIVE LIST OF ALGEBRA 2 FORMULAS MATTERS

Algebra 2 is a pivotal course in high school mathematics, bridging foundational concepts from Algebra 1 and Geometry to the more abstract thinking required for precalculus and calculus. At its core, Algebra 2 introduces students to polynomial functions, exponential

and logarithmic relationships, complex numbers, and data analysis. Mastering the material often comes down to memorizing and knowing how to apply key formulas. A well-organized **list of algebra 2 formulas** can serve as a lifeline for homework, test preparation, and real-world problem solving. Whether you are a student reviewing for finals, a teacher creating a reference sheet, or someone returning to math after many years, having all those essential equations in one place saves time and

reduces frustration. This article provides a comprehensive, clearly categorized guide to the most important formulas you will encounter in a typical Algebra 2 course. We will explore each formula's purpose, its components, and when to use it, so you can build both fluency and confidence.

LINEAR FUNCTIONS AND EQUATIONS

Linear relationships are the foundation of algebra. Even as Algebra 2 dives into higher-degree polynomials, working with lines remains essential for modeling and for understanding slopes.

Slope Formula

Given two points $((x_1, y_1))$ and $((x_2, y_2))$:

- **Slope (m)** =
$$\frac{y_2 - y_1}{x_2 - x_1}$$

This formula expresses the rate of change of a linear function. It is used to determine whether lines are parallel ($(m_1 = m_2)$) or perpendicular ($(m_1 \cdot m_2 = -1)$).

Forms of Linear Equations

- **Slope-intercept form:** $(y = mx + b)$

- **Point-slope**

form: $(y - y_1 = m(x - x_1))$

- **Standard form:**

$(Ax + By = C)$

Each form has advantages: slope-intercept is ideal for graphing; point-slope is perfect when you know a point and the slope; standard form is useful for quickly finding intercepts.

Distance and Midpoint Formulas

- **Distance:** $(d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2})$

- **Midpoint:**

$(\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right))$

$(\frac{y_1 + y_2}{2}, \frac{x_1 + x_2}{2})$

These formulas extend the coordinate geometry skills from Algebra 1 into more complex problem contexts in Algebra 2, such as finding lengths of segments or centers of figures.

QUADRATIC FUNCTIONS AND EQUATIONS

Quadratic equations are central to Algebra 2. They appear in projectile motion, area optimization, and many science applications.

Standard Form of a Quadratic

$(f(x) = ax^2 + bx + c)$, where

$$(a \neq 0)$$

Vertex Form

- $f(x) = a(x - h)^2 + k$
- The vertex is at (h, k) .

Quadratic Formula (for solving $ax^2 + bx + c = 0$)

- $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

The discriminant $(b^2 - 4ac)$ tells us about the nature of roots: positive means two real solutions, zero means one real solution, negative means two complex solutions.

Factoring Formulas

- **Difference of squares:** $a^2 - b^2 = (a - b)(a + b)$
- **Perfect square trinomial:** $a^2 \pm 2ab + b^2 = (a \pm b)^2$

Recognizing these patterns helps solve quadratic equations efficiently without always using the quadratic formula.

POLYNOMIAL

FUNCTIONS

Algebra 2 expands polynomial operations to include higher degrees. The following formulas are essential for factoring and analyzing polynomial behavior.

Sum and Difference of Cubes

- $a^3 + b^3 = (a + b)(a^2 - ab + b^2)$
- $a^3 - b^3 = (a - b)(a^2 + ab + b^2)$

Remainder

er and Factor Theorems

- **Remainder Theorem:** If polynomial $P(x)$ is divided by $(x - c)$, the remainder is $P(c)$.
- **Factor Theorem:** $(x - c)$ is a factor of $P(x)$ if and only if $P(c) = 0$.

Rational Root Theorem

For a polynomial $a_n x^n + a_{n-1} x^{n-1} + \dots +$

a_0), possible rational roots are $\frac{\text{factors of } a_0}{\text{factors of } a_n}$.

Fundamental

Theorem of

Algebra

Every nonzero polynomial of degree n has exactly n complex roots (counting multiplicities).

EXPONENTIAL AND LOGARITHMIC FUNCTIONS

These functions model growth, decay, and many natural processes. A firm grasp

of their formulas is critical for Algebra 2 success.

Laws of Exponents

- $a^m \cdot a^n = a^{m+n}$
- $\frac{a^m}{a^n} = a^{m-n}$
- $(a^m)^n = a^{mn}$
- $a^{-n} = \frac{1}{a^n}$
- $a^{1/n} = \sqrt[n]{a}$

Definition of a Logarithm

m

- $\log_b(x) = y$ if and only if $b^y = x$

Logarithmic Properties

- $\log_b(MN) = \log_b M + \log_b N$
- $\log_b\left(\frac{M}{N}\right) = \log_b M - \log_b N$
- $\log_b(M^p) = p \log_b M$
- $\log_b(1) = 0$
- $\log_b(b) = 1$
- **Change-of-base formula:** $\log_b x = \frac{\log_a x}{\log_a b}$

Exponential and Logarithmic Equations

When solving equations, these formulas allow you to move between exponential and logarithmic forms. Common applications include compound interest $A = Pe^{rt}$ (continuous) or $A = P(1 + r/n)^{nt}$.

RADICAL EXPRESSIONS AND COMPLEX NUMBERS

Radicals and imaginary numbers often appear

together. The following list of algebra 2 formulas helps simplify radicals and operate with complex numbers.

Simplifyin

g

Radicals

- $\sqrt[n]{a^n} = |a|$ if n is even, or just a if n is odd.
- $\sqrt[n]{ab} = \sqrt[n]{a} \cdot \sqrt[n]{b}$
- $\sqrt[n]{\frac{a}{b}} = \frac{\sqrt[n]{a}}{\sqrt[n]{b}}$

Imaginar y Unit

and Complex Numbers

- $i = \sqrt{-1}$
- $i^2 = -1$
- $i^3 = -i$
- $i^4 = 1$ (cyclic pattern)
- **Complex number form:** $a + bi$ where a and b are real numbers.
- **Conjugate:** $a + bi$ and $a - bi$
- **Modulus:** $|a + bi| = \sqrt{a^2 + b^2}$

RATIONAL FUNCTIONS AND EQUATIONS

Rational functions involve fractions with polynomials in the numerator and

denominator. Working with these requires careful algebraic manipulation.

Simplifying Rational Expressions

Factor numerator and denominator completely, then cancel common factors. Keep track of excluded values where denominator equals zero.

Solving Rational

Equations

Multiply both sides by the least common denominator (LCD) to clear fractions, then solve the resulting polynomial equation. Always check for extraneous solutions.

Asymptotes

- **Vertical asymptote:**
Values that make denominator zero (after canceling common factors).
- **Horizontal asymptote:**
Compare degrees: if numerator degree < denominator degree, $y=0$; if equal, $y = \text{ratio}$

of leading coefficients; if greater, no horizontal asymptote (oblique possible).

SEQUENCES AND SERIES

Algebra 2 introduces arithmetic and geometric sequences, leading to series and summation notation.

Arithmeti C Sequence S

- $(a_n = a_1 + (n-1)d)$
- Sum of first (n) terms: $(S_n = \frac{n}{2}(a_1 + a_n))$ or $(S_n$

$$= \frac{n}{2}[2a_1 + (n-1)d]$$

Geometri C Sequence S

- $(a_n = a_1 \cdot r^{n-1})$
- Sum of first (n) terms: $(S_n = a_1 \frac{1 - r^n}{1 - r})$ for $(r \neq 1)$
- Infinite geometric series (if $(|r| < 1)$): $(S_{\infty} = \frac{a_1}{1 - r})$

Summati

on

Notation

$\sum_{k=1}^n a_k$ is a compact way to write a series. Knowing the formulas above makes evaluating sums straightforward.

PROBABILITY AND COMBINATORICS

Counting and probability are integral parts of the Algebra 2 curriculum, especially for data analysis.

Factorial

- $n! = n \cdot (n-1) \cdot (n-2) \cdots 1$

Permutati

ons

(order

matters)

- ${}_nP_r = \frac{n!}{(n-r)!}$

Combinat

ions

(order

does not

matter)

- ${}_nC_r = \frac{n!}{r!(n-r)!}$

Binomial Theorem

$$\left((x + y)^n = \sum_{k=0}^n \binom{n}{k} x^{n-k} y^k \right)$$